



[cancer.org](https://www.cancer.org) | 1.800.227.2345

If You Have Bladder Cancer

- [What is bladder cancer?](#)
- [How does the doctor know I have bladder cancer?](#)
- [How serious is my cancer?](#)
- [What kind of treatment will I need?](#)
- [What will happen after treatment?](#)

What is bladder cancer?

Cancer can start any place in the body. Cancer that starts in the bladder is called **bladder cancer**. It starts when cells in the bladder grow out of control and crowd out normal cells. This makes it hard for the body to work the way it should.

Bladder cancer cells can sometimes spread to other parts of the body, such as the bones, and grow there. This spread is called **metastasis**

The "transitional epithelium" in the picture below is the lining layer where most bladder cancers start. Over time they can spread deeper into the other layers.



Ask your doctor to use this picture to show you where the cancer is.

Questions to ask the doctor

- Why do you think I have bladder cancer?
- Is there a chance I don't have cancer?
- Would you please write down the kind of cancer you think I might have?

[Signs and symptoms of bladder cancer](#)¹ are problems peeing, pain when peeing, needing to go more often than normal, and seeing blood in your urine.

If signs suggest you might have bladder cancer, exams and tests will be done. Here are some of the [tests you may need](#)²:

Physical exam: The doctor will check you for signs of bladder cancer and other health problems. This might include a rectal exam, during which a gloved finger is put into your rectum. If you are a woman, a pelvic exam might also be done. During these exams, the doctor can sometimes feel a bladder tumor.

Urine tests: For these tests, you'll be asked to pee in a cup. Your urine is then tested for cancer cells, blood, or certain proteins (called biomarkers or tumor markers).

Cystoscopy: For this exam, a doctor called a urologist looks at the inside of your bladder using a tool called a cystoscope. This is a thin tube with a tiny light and camera on its end. It's put through the opening of your urethra and moved up into your bladder. This exam can also be used to remove (biopsy) small pieces of the bladder wall, which can then be tested for cancer. Saltwater washings of the inside of your bladder may also be collected to look for cancer cells.

Blue light cystoscopy: Sometimes, special drugs are put into the bladder during the exam. Cancer cells soak up these drugs and then glow when the doctor shines a blue light through the scope. This can help the doctor see cancer cells that might have been missed with the normal light.

TURBT: In this procedure, a rigid cystoscope called a resectoscope is put into the bladder. This lets the doctor remove tumors as well as some of the muscle layer of the bladder wall, which can then be checked in the lab for cancer cells. A TURBT can also be a form of treatment for early bladder cancers.

Lab tests of biopsy samples: Any samples removed during either a cystoscopy or a TURBT are tested in the lab for cancer cells. This is needed to know for sure if you have bladder cancer. Other tests may be done on the cells to learn more about the cancer.

CT scan: This is sometimes called a CAT scan. It uses x-rays to make detailed pictures of the inside of your body. It can show if the cancer has spread outside the bladder.

MRI: MRIs use radio waves and strong magnets instead of x-rays to take detailed pictures. MRIs can be very helpful in finding cancer that has spread outside the bladder.

Ultrasound: This test uses sound waves to make pictures of the organs inside your

body, like your bladder and kidneys. It can help show the size of a bladder cancer and if it has spread.

X-ray tests: These tests aren't usually done, because other tests such as CT or MRI scans are often all that's needed. For one type of x-ray test (called an IVP), a dye is put

bladder wall, and possibly deeper. These cancers are more likely to spread, and they tend to be harder to treat.

Grade

The grade refers to how the cancer looks under the microscope.

- **Low-grade** bladder cancers look a lot like normal bladder cells. They tend to grow and spread slowly.
- **High-grade** bladder cancers look less like normal bladder cells. These cancers are more likely to grow and spread. They can be harder to treat.

Gene and protein changes

Lab tests may be done to check the cancer cells for certain gene or protein changes (sometimes called **biomarkers**). This might affect your treatment options.

Stage

Your doctor will also want to find out the [stage of your cancer](#)³ to help decide what type of treatment is best for you. The stage describes the growth or spread of the cancer in the bladder, as well as if it has spread to other parts of your body.

Your cancer can be stage 0, 1, 2, 3, or 4. The lower the number, the less the cancer has spread. A higher number, like stage 4, means a more serious cancer that has spread outside the bladder. Ask your doctor about the cancer stage and what it means for you.

Risk groups

If you have NMIBC, the cancer might also be put into a risk group (low, intermediate, or high), based on certain factors. This can give doctors an idea of how likely it is that the cancer will come back or keep growing, which might affect your treatment options.

Questions to ask the doctor

- Has the cancer invaded the muscle layer of the bladder?
- Do you know the stage and grade of the cancer?

- If not, how and when will you find out the stage and grade?
- Would you explain what these mean in my case?
- What will happen next?

What kind of treatment will I need?

There are many [ways to treat bladder cancer](#)⁴. You might want to get a [second opinion](#)⁵ about the best treatment plan for you.

Bladder cancer is most often treated with:

- Surgery
- Intravesical treatment
- Radiation
- Chemotherapy (chemo)
- Immune therapy
- Targeted drugs

Sometimes more than one type is used. The treatment plan that's best for you depends on:

- The stage and grade of the cancer
- How far the cancer has spread into the bladder wall
- The chance that a type of treatment will cure the cancer or help in some way
- Any other health problems you have
- Your feelings about the treatment and the side effects that come with it

Surgery for bladder cancer

Surgery is done for most bladder cancers. The type you have depends on the stage of the cancer and other factors, such as your health and preferences.

Removing the tumor from the inside bladder (a transurethral resection of bladder tumor, or TURBT) is the most common surgery for early bladder cancer. This can be done during a [cystoscopy](#)⁶.

When the cancer is more invasive, the cancer is removed along with part of the bladder

(partial cystectomy) or the entire bladder (radical cystectomy).

If only part of the bladder is removed, you'll still be able to hold and release urine as normal, though in smaller amounts. If the entire bladder is removed, you'll need another way to store and pass urine. Your doctor can explain the options for this.

Side effects of surgery

Any type of surgery can have some [risks and side effects](#)⁷. For instance, removing the bladder not only changes how your body passes urine, but it can also cause sexual side effects. If you have these or any other problems, let your doctors know. There are ways to help deal with many side effects.

Intravesical treatment

For this treatment, a drug is put inside the bladder. This might be a chemo drug or a drug such as BCG that boosts the immune system. This is often done after surgery in people who still have their bladder. It can help lower the chance that the cancer will come back in the bladder.

The drug affects the cells lining the inside of the bladder, but it has little to no effect on cells in other places. This can help limit side effects. But this also means it can't treat invasive bladder cancers or those that have spread outside the bladder.

Side effects of intravesical treatment

Some people have a burning feeling in their bladder after treatment. Other side effects depend on which drug is used. Some side effects may feel like having the flu, such as fever, chills, and feeling tired.

Chemo

Chemo is the use of drugs to kill cancer cells. The drugs may be given into a vein or taken as pills. They go into the blood and spread through the body.

For early-stage bladder cancers, chemo may be used:

- Before surgery to shrink a tumor
- After surgery to kill any cancer cells that remain
- With radiation to help it work better

- As a treatment for advanced bladder cancers, such as those that have spread to other parts of the body

Chemo is given in cycles or rounds. Each round of treatment is followed by a break. Most of the time, 2 or more chemo drugs are given. Treatment often lasts for many months.

Side effects of chemo

Chemo can make you feel very tired, sick to your stomach, and cause your hair to fall out. But these problems go away after treatment ends.

There are ways to treat most [chemo side effects](#)⁸. If you have side effects, talk to your cancer care team so they can help.

Radiation treatment

In radiation therapy, a machine sends x-rays to a specific part of the body to kill cancer cells.

Radiation treatment for bladder cancer can be used:

- To treat early-stage cancer after surgery
- As the main treatment for early-stage cancer if you can't have (or don't want) surgery
- As part of the treatment for advanced bladder cancer

Radiation is often given along with chemo. Certain chemo drugs can help the radiation work better.

Side effects of radiation treatments

The most common side effects of radiation to the bladder are:

- Feeling tired
- Easy bruising or bleeding
- Increased risk of infection

Most side effects get better after treatment ends. Some might last longer. Talk to your doctor about what you can expect.

Immunotherapy

Immunotherapy is treatment that boosts your immune system to attack the cancer cells. Different types of immunotherapy can be used to treat bladder cancer. These drugs can either be put right into the bladder (as a liquid) to treat some early-stage bladder cancers, or they can be given into a vein to help treat more advanced bladder cancers.

Side effects of immunotherapy

Immunotherapy can cause different side effects depending on which drug is used. Most side effects tend to be mild, such as feeling tired or having an upset stomach, but in some people the side effects can be serious. Most side effects go away after treatment ends.

If you have side effects, talk to your cancer care team so they can help.

Targeted drugs

your doctor if your clinic or hospital conducts clinical trials. See [Clinical Trials](#)⁹ to learn more.

What about other treatments that I hear about?

When you have bladder cancer, you might hear about other ways to treat cancer or treat your symptoms that are not standard medical treatments. These treatments can be vitamins, herbs, special diets, and other things. You may be curious about these treatments.

Some of these are known to help, but many have not been tested. Some have been shown not to help. A few have even been found to be harmful. Talk to your doctor about anything you're thinking about using, whether it's a vitamin, a diet, or anything else.

Questions to ask the doctor

- What are my treatment options?
 - What treatment do you think is best for me? Why?
 - What's the goal of this treatment? Do you think it could cure the cancer?
 - Will treatment include surgery? If so, who will do the surgery?
 - What will the surgery be like?
 - Does my bladder need to be removed? If so, how will I pee after surgery?
 - Will I have other types of treatment?
 - What's the goal of these treatments?
 - What should I do to be ready for treatment?
 - Is there anything I can do to help the treatment work better?
- What side effects c95.35 39measleg521ld0fter surgery?

second bladder cancer.

If you have no signs of cancer, most experts advise seeing your doctor every 3 to 6 months. These visits might include urine tests, blood work, and other tests. If you still have your bladder, you will need regular exams of your bladder, too. The time between doctor visits may be longer after a few years if no new cancers are seen.

Having cancer and dealing with treatment can be hard, but it can also be a time to look at your life in new ways. You might be thinking about how to improve your health. Call us or talk to your doctor to find out what you can do to feel better.

You can't change the fact that you have cancer. What you can change is how you live the rest of your life, making healthy choices and feeling as well as you can.

[For connecting and sharing during a cancer journey](#)

Anyone with cancer, their caregivers, families, and friends, can benefit from help and support. The American Cancer Society offers the Cancer Survivors Network (CSN), a safe place to connect with others who share similar interests and experiences. We also partner with CaringBridge, a free online tool connecting people with similar experiences.

9. www.cancer.org/cancer/managing-cancer/making-treatment-decisions/clinical-trials.html
10. www.cancer.org/cancer/types/bladder-cancer/after-treatment/follow-up.html
11. www.cancer.org

Words to know

Bladder: A hollow organ that stores urine.

Biopsy (BY-op-see): Taking out tiny pieces of tissue to see if they contain cancer cells.

Cystectomy (sis-TEK-tuh-mee): Surgery to take out the bladder.

Cystoscopy (sis-TAH-scuH-pee): A procedure to look at the inside of the bladder with a thin, lighted tube called a cystoscope (SIS-tah-scope).

Intravesical (in-truh-VESS-uh-cull): Within or inside the bladder.

Invasive cancer: Cancer that has spread beyond the layer of cells where it first began.

Kidneys: A pair of bean-shaped organs, each about the size of a fist, that make urine.

Lymph nodes (limf nodes): Small bean-sized parts of the immune system found all over the body and connected by lymph vessels.

Metastasis (meh-TAS-tuh-sis): The spread of cancer cells from where they started to other places in the body.

Non-invasive cancer: Cancer that's still only in the layer of cells where it first began.

Partial cystectomy (sis-TEK-tuh-mee): Surgery to remove only part of the bladder.

Radical cystectomy: Surgery to remove the entire bladder.

TURBT: A procedure that removes small pieces from the inside of the bladder to see if someone has bladder cancer. It can also be used to treat some small bladder cancers.

Ureter (YUR-uh-tur): A tube that carries urine from each kidney to the bladder.

Urethra (yur-EETH-ruh): The tube that carries urine from the bladder to the outside of the body.

Urine: Liquid waste made by the kidneys and stored in the bladder. Also called pee.

Urologist (yur-OL-uh-jist): A doctor who's an expert in treating problems of the urinary tract, as well as the genital area in men.

How can I learn more?

We have a lot more information for you. You can find it online at www.cancer.org¹¹. Or, you can call our toll-free number at 1-800-227-2345 to talk to one of our cancer information specialists.

Last Revised: March 12, 2024

Written by

The American Cancer Society medical and editorial content team
(<https://www.cancer.org/cancer/acs-medical-content-and-news-staff.html>)

Our team is made up of doctors and oncology certified nurses with deep knowledge of cancer care as well as editors and translators with extensive experience in medical writing.

American Cancer Society medical information is copyrighted material. For reprint requests, please see our Content Usage Policy (www.cancer.org/about-us/policies/content-usage.html).

cancer.org | 1.800.227.2345