



cancer.org | 1.800.227.2345

Mesothelioma Causes, Risk Factors, and Prevention

Learn about the risk factors for mesothelioma and what you might be able to do to help lower your risk.

Risk Factors

A risk factor is anything that affects your chance of getting a disease such as cancer. Learn more about the risk factors for mesothelioma.

- [Risk Factors for Mesothelioma](#)
- [What Causes Mesothelioma?](#)

Prevention

There's no way to completely prevent mesothelioma. But there are things you can do that might lower your risk. Learn more.

- [Can Mesothelioma Be Prevented?](#)

Risk Factors for Mesothelioma

- [Asbestos](#)
- [Zeolites](#)

- [Radiation](#)
- [SV40 virus](#)
- [Age](#)
- [Sex](#)
- [Gene changes](#)

A risk factor is anything that increases your chance of getting a disease such as cancer. Different cancers have different risk factors. Some risk factors, like smoking, can be changed. Others, like a person's age or family history, can't be changed. But having a known risk factor, or even many, does not mean that you will get the disease. And some people who get the disease may have few or no known risk factors. Researchers have found some factors that increase a person's risk of mesothelioma.

Asbestos

The main risk factor for **pleural mesothelioma** is exposure to asbestos. In fact, most cases of pleural mesothelioma have been linked to high levels of asbestos exposure, usually in the workplace.

Asbestos is a group of minerals that occur naturally as bundles of tiny fibers. These fibers are found in soil and rocks in many parts of the world.

When asbestos fibers in the air are inhaled, they can get into the lungs. Fibers that stay in the lungs can travel to the ends of the small airways and enter the pleural lining of the lung and chest wall. These fibers can then injure the cells of the pleura, and, over time, cause mesothelioma. Asbestos fibers can also damage cells of the lung and result in **asbestosis** (scar tissue in the lung) and/or [lung cancer](#)¹ .

Peritoneal mesothelioma can form in the abdomen when inhaled asbestos fibers are coughed up and then swallowed.

Many people are exposed to very low levels of naturally occurring asbestos in outdoor air. It's in dust that comes from rocks and soil containing asbestos. This is more likely to happen in areas where rocks have higher asbestos content. In some areas, asbestos can be found in the water supply as well as in the air.

In the past, asbestos was used in many products because it was heat and fire-resistant. The link between asbestos and mesothelioma is now well known, and most of its use in the United States stopped several decades ago, but it's still used in some products.

Still, millions of Americans may already have been exposed to asbestos. People at risk

treated with radiation, this cancer is still rare in these patients.

SV40 virus

Some studies have raised the possibility that infection with simian virus 40 (SV40) might increase the risk of developing mesothelioma. But most experts agree that at this time we still don't know if SV40 is responsible for some mesotheliomas. This important topic is still being researched.

Age

The risk of mesothelioma increases with age. Mesothelioma can occur in young people (even children), but it's rare in people under age 45. About 2 out of 3 people with mesothelioma of the chest are 65 or older.

Sex

Mesothelioma is much more common in men than in women. This is probably because men have been more likely to work in jobs with heavy exposure to asbestos.

Gene changes

A mutation or change in the gene called *BAP1* can be passed in families and has been linked to mesothelioma. But *BAP1* mutations are rare.

Hyperlinks

1. www.cancer.org/cancer/types/lung-cancer.html
2. www.cancer.org/cancer/risk-prevention/chemicals/asbestos.html

References

American Society of Clinical Oncology. Mesothelioma: Risk Factors. 07/2017. Accessed at www.cancer.net/cancer-types/mesothelioma/risk-factors on October 17, 2018.

Hung YP, Chirieac LR. Novel insights and recent discoveries on the genetics and pathogenesis of malignant mesothelioma. *J Thorac Dis.* 2018;10(3):1314-1317.

Mesothelioma Applied Research Foundation. The Debate Surrounding SV40 and Its Role in the Development of Mesothelioma. Accessed at www.curemeso.org/understanding-mesothelioma/mesothelioma-development/mesothelioma-role-simian-virus-40-sv40/ on October 17, 2018.

National Comprehensive Cancer Network, Clinical Practice Guidelines in Oncology (NCCN Guidelines®), Malignant Pleural Mesothelioma, Version 2.2018 -- February 26, 2018. Accessed at www.nccn.org/professionals/physician_gls/pdf/mpm.pdf on October 17, 2018.

Last Revised: November 16, 2018

What Causes Mesothelioma?

Researchers have found several factors that increase a person's risk of mesothelioma, but it's not yet clear exactly how all of these factors might cause this cancer.

Cancers, including mesotheliomas, occur when the DNA in cells is damaged. DNA is the chemical in each of our cells that makes up our *genes* – the instructions for how our cells function. We usually look like our parents because they are the source of our DNA. But DNA affects more than how we look. Some genes control when cells in the body grow, divide into new cells, and die. Changes in these genes may cause cells to grow out of control, which can lead to cancer.

Asbestos exposure is the main cause of pleural mesothelioma. About 8 out of 10 people with mesothelioma have been exposed to asbestos. When asbestos fibers are breathed in, they travel to the ends of small air passages and reach the pleura, where they can cause inflammation and scarring. This may damage cells' DNA and cause changes that result in uncontrolled cell growth. If swallowed, these fibers can reach the abdominal lining, where they can have a role in causing peritoneal mesothelioma. But most people exposed to asbestos, even in large amounts, do not get mesothelioma.

Radiation treatments for other cancers have been linked to mesothelioma in some studies. Radiation can damage the cells' DNA, leading to out-of-control cell growth.

References

containing asbestos generally do not pose a health risk. They may pose a risk if they are damaged, disturbed in some way, or deteriorate over time and release asbestos fibers into the air. By federal law, all schools are required to inspect materials with asbestos regularly and must have a plan in place for managing them.

Hyperlinks

1. www.cancer.org/cancer/risk-prevention/chemicals/asbestos.html

References

Mesothelioma Cancer Alliance. What Is Asbestos? Accessed at www.mesothelioma.com/asbestos-exposure/what-is-asbestos.htm on October 17, 2018.

Last Revised: November 16, 2018

Written by

The American Cancer Society medical and editorial content team
(<https://www.cancer.org/cancer/acs-medical-content-and-news-staff.html>)

Our team is made up of doctors and oncology certified nurses with deep knowledge of cancer care as well as editors and translators with extensive experience in medical writing.

American Cancer Society medical information is copyrighted material. For reprint requests, please see our Content Usage Policy (www.cancer.org/about-us/policies/content-usage.html).

cancer.org | 1.800.227.2345