

cancer.org | 1.800.227.2345

Oral Cavity and Oropharyngeal Cancer Causes, Risk Factors, and Prevention

Learn about the risk factors for cancers of the oral cavity (mouth cancer) and oropharynx (throat cancer) and what you might be able to do to help lower your risk.

Risk Factors

A risk factor is anything that increases your chance of getting a disease such as cancer. Learn more about the risk factors for oral cavity and oropharyngeal cancer.

- Risk Factors for Oral Cavity and Oropharyngeal Cancers
- What Causes Oral Cavity and Oropharyngeal Cancers?

Prevention

There's no way to prevent all oral cavity and oropharyngeal cancers. But there are things you can do that might help lower your risk. Learn more here.

Can Oral Cavity and Oropharyngeal Cancers Be Prevented?

Risk Factors for Oral Cavity and Oropharyngeal Cancers

- Tobacco and alcohol use
- Human papillomavirus (HPV) infection
- Sex
- Excess body weight
- Age
- Ultraviolet (UV) light

products. People who continue to smoke while getting cancer treatment can have trouble with wound healing, more side effects from radiation therapy or chemotherapy, a higher risk of infection, and worse outcomes. Also, continuing to smoke and use oral tobacco products after cancer treatment, greatly increases the risk of developing a second cancer² in the mouth, throat, larynx (voice box), lung and other organs.

If you are thinking about quitting tobacco and need help, call the American Cancer Society at 1-800-227-2345. A tobacco cessation and counseling program can help increase your chances of quitting for good. See How to Quit Smoking or Smokeless Tobacco for more information.

Drinking alcohol increases the risk of developing oral cavity and oropharyngeal cancers. Heavy drinkers have a higher risk than light drinkers.

Smoking and drinking alcohol together multiplies the risk of these cancers. The risk of these cancers in people who drink and smoke heavily is about 30 times higher than the risk in people who don't smoke or drink.

Betel quid and gutka

In Southeast Asia, South Asia, and certain other areas of the world, many people chew betel quid, which is made up of areca nut (betel nut), spices, lime, and other ingredients. Many people in these areas also chew gutka, a mixture of betel quid and tobacco. People who chew betel quid or gutka have an increased risk of cancer of the mouth.

Human papillomavirus (HPV) infection

Human papillomavirus (HPV) is a group of more than 150 types of viruses. They're calledformation.

outcomes (prognoses) than tumors not caused by HPV because chemotherapy and radiation treatments work better for these cancers. This improvement in outcome

Genetic syndromes

People with syndromes caused by <u>inherited defects (mutations) in certain genes</u>⁶ have a very high risk of mouth and middle throat cancer.

- Fanconi anemia: People with this syndrome often have blood problems at an early age, which may lead to leukemia or myelodysplastic syndrome. They also have a very high risk of cancers of the mouth and throat.
- Dyskeratosis congenita: This is a genetic syndrome that can cause aplastic anemia, skin rashes, and abnormal fingernails and toenails. People with this syndrome also have a very high risk of developing head and neck cancers, especially of the mouth and throat, at a young age.

Unproven or controversial risk factors

Mouthwash

Some studies have suggested that mouthwash with a high alcohol content might be linked to a higher risk of oral and oropharyngeal cancers. But recent research has questioned these results. Studying this possible link is complicated by the fact that people who smoke and frequent drinkers (who already have an increased risk of these cancers) are more likely to use mouthwash than people who neither smoke nor drink.

syclikel0.78F2 12 Tf 0 0 0 rg /GS156 gs6(especIt'sj 0 costed that moutlita-term ihs361.55le mout

Clin Epigenetics. 2017;9:124.

Chainani-Wu N. Diet and oral, pharyngeal, and esophageal cancer. *Nutr Cancer*. 2002;44(2):104-126. doi:10.1207/S15327914NC4402_01.

Cohen N, Fedewa S, Chen AY. Epidemiology and Demographics of the Head and Neck Cancer Population. *Oral Maxillofac Surg Clin North Am.* 2018;30(4):381-395. doi:10.1016/j.coms.2018.06.001.

pharyngeal cancer risk. *Oral Oncol.* 2009;45(6):461-467. doi:10.1016/j.oraloncology.2008.09.002.

Reidy JT, McHugh EE, Stassen LF. A review of the role of alcohol in the pathogenesis of oral cancer and the link between alcohol-containing mouthrinses and oral cancer. *J Ir Dent Assoc.* 2011;57(4):200-202.

Singhvi HR, Malik A, Chaturvedi P. The Role of Chronic Mucosal Trauma in Oral Cancer: A Review of Literature. *Indian J Med Paediatr Oncol.* 2017;38(1):44-50. Doi:10.4103/0971-5851.203510.

Shah A, Malik A, Garg A, et al. Oral sex and human papilloma virus-related head and neck squamous cell cancer: a review of the literature. *Postgrad Med J*. 2017;93(1105):704-709.

What Causes Oral Cavity and

Oropharyngeal Cancers?

- Tobacco and alcohol and oral or oropharyngeal cancers
- Human papillomavirus (HPV) and oral or oropharyngeal cancers
- Inherited or acquired gene mutations and oral or oropharyngeal cancers

Doctors and scientists can't say for sure what causes each case of oral cavity or oropharyngeal cancer. But they do know many of the <u>risk factors</u> and how some of them may lead to cells becoming cancer.

For human cells to be made, this mostly depends on the information in the cells' DNA. DNA is the chemical in our cells that makes up our genes, which control how our cells work. We look like our parents because they are the source of our DNA. But DNA affects more than just how we look.

Some genes control when cells grow, divide, and die:

- Genes that help cells normally grow, divide, and stay alive are called **proto-oncogenes**. When a proto-oncogene mutates (changes), the gene becomes abnormal and is then called an **oncogene**.
- Genes that help keep cell growth under control or make cells die at the right time are called tumor suppressor genes.

Cancers can be caused by DNA mutations (gene changes) that turn on oncogenes or turn off tumor suppressor genes. This leads to cells growing out of control. Changes in many different genes are usually needed to cause oral cavity and oropharyngeal cancer.

For more about how genes changes can lead to cancer, see <u>Genes and Cancer</u>¹.

Tobacco and alcohol and oral or oropharyngeal cancers

Scientists believe that some risk factors, such as tobacco or heavy alcohol use, may cause these cancers by damaging the DNA of cells that line the inside of the mouth and throat.

When tobacco and alcohol damage the cells lining the mouth and throat, the cells in this layer need to divide more often and make more copies of themselves. This increases the chance to make mistakes when copying their DNA, which may increase the

possibility of becoming cancer.

Many of the chemicals found in tobacco can damage DNA directly. Scientists are not sure whether alcohol directly damages DNA, but they have shown that alcohol helps many DNA-damaging chemicals get into cells more easily. This may be why the combination of tobacco and alcohol damages DNA far more than tobacco alone. Acetaldehyde, a breakdown product of alcohol, is found in saliva (spit). It has been shown to get in the way of normal DNA repair, as well as disturb other functions of DNA. The level of acetaldehyde goes up as people drink more alcohol which may be one way alcohol damages the cells lining the oral cavity.

This damage can cause certain genes (for example, those in charge of starting or stopping cell growth) to not work well. Abnormal cells can begin to grow out of control, forming a tumor.

Human papillomavirus (HPV) and oral or oropharyngeal cancers

Certain types of <u>human papillomavirus</u>² (HPV) infections (mainly HPV16), are important causes of most oropharyngeal (middle throat) cancers. HPV makes 2 proteins known as E6 and E7 which turn off some tumor suppressor genes, such as Rb and p53. In the throat, too much of the E6 and E7 proteins are made. This may allow the cells lining the oropharynx to grow out of control and to develop changes in more genes, which in some cases can lead to cancer.

The number of oropharyngeal cancers caused by HPV infections is going up. They tend to be found in younger people who don't smoke or drink alcohol, and tend to have a better outcome (prognosis) than throat cancers that are not caused by HPV.

Inherited or acquired gene mutations and oral or oropharyngeal cancers

Some people inherit DNA mutations (changes) from their parents that increase their risk

chemicals, like those found in tobacco smoke. Several different gene changes are probably needed for cancer to develop, and not all of these changes are understood at this time.

Hyperlinks

- 1. www.cancer.org/cancer/understanding-cancer/genes-and-cancer.html
- 2. www.cancer.org/cancer/risk-prevention/hpv.html

References

Basen-Engquist K, Borwn P, Coletta AM, Savage M, Laresso KC, Hawk E. Ch. 22 - Lifestyle and Cancer Prevention. In: Niederhuber JE, Armitage JO, Doroshow JH, Kastan MB, Tepper JE, eds. *Abeloff's Clinical Oncology*. 6th ed. Philadelphia, Pa. Elsevier; 2020.

Leeman JE, Katabi N, Wong RJ, Lee NY, Romesser PB. Ch. 65 - Cancer of the Head and Neck. In: Niederhuber JE, Armitage JO, Doroshow JH, Kastan MB, Tepper JE, eds. *Abeloff's Clinical Oncology*. 6th ed. Philadelphia, Pa. Elsevier; 2020.

Mendenhall WM, Dziegielewski PT, Pfister DG. Chapter 45- Cancer of the Head and Neck. In: DeVita VT, Lawrence TS, Rosenberg SA, eds. *DeVita, Hellman, and Rosenberg's Cancer: Principles and Practice of Oncology.* 11th ed. Philadelphia, Pa: Lippincott Williams & Wilkins; 2019.

Tumban E. A Current Update on Human Papillomavirus-Associated Head and Neck Cancers. *Viruses*. 2019;11(10):922. Published 2019 Oct 9. doi:10.3390/v11100922.

Last Revised: March 23, 2021

Can Oral Cavity and Oropharyngeal Cancers Be Prevented?

- Avoid tobacco and alcohol
- Avoid HPV infection/Get the HPV vaccine
- Limit exposure to ultraviolet (UV) light
- Maintain a healthy weight and eating pattern
- Get regular dental checkups

Not all cases of oral cavity (mouth) and oropharyngeal (middle throat) cancer can be prevented, but the risk of developing these cancers can be greatly reduced if you take steps to avoid certain risk factors.

Avoid tobacco and alcohol

Use of tobacco and alcohol are among the most important risk factors for these cancers. Not starting to smoke or use oral tobacco products is the best way to limit the risk of getting these cancers. Quitting tobacco¹ also greatly lowers your risk of developing these cancers, even after many years of use. Heavy alcohol use is a risk factor on its own. It also greatly increases the cancer-causing effect of tobacco. So it's especially important to avoid smoking and alcohol as well as the combination of alcohol and smoking.

Avoid HPV infection/Get the HPV vaccine

The risk of human papillomavirus (HPV) infection of the mouth and middle throat is increased in those who have oral sex and multiple sex partners. These infections are also more common in people who smoke, which may be because the smoke damages their immune system or the cells that line the mouth and throat.

HPV is very common and rarely causes symptoms. And even though HPV infection is linked to most cases of oropharyngeal cancer, most people with HPV infections of the

See HPV³ to learn more.

Limit exposure to ultraviolet (UV) light

<u>Ultraviolet radiation</u>⁴ is an important and avoidable risk factor for cancer of the lips, as well as for <u>skin cancer</u>⁵. If possible, limit the time you spend outdoors during the middle of the day, when the sun's UV rays are strongest. If you are out in the sun, wear a widebrimmed hat and use sunscreen and lip balm with a sun protection factor (SPF) of at least 30.

Maintain a healthy weight and eating pattern

Poor nutrition and excess body weight have been linked to oral cavity and oropharyngeal cancers. Following a healthy eating pattern with more plant-based foods, such as non-starchy vegetables and whole fruit may help lower your risk of these cancers (and many others).

The American Cancer Society recommends following a healthy eating pattern that includes plenty of fruits, vegetables, and whole grains, and that limits or avoids red and processed meats, sugary drinks, and highly processed foods. In general, eating a healthy diet is much better than adding vitamin supplements to an otherwise unhealthy diet. See the American Cancer Society Guidelines for Diet and Physical Activity for Cancer Prevention⁶ for our full guidelines.

Get regular dental checkups9 gs ()Tj 0 g 0 0 0 rg /GS500 gs (for our full g (in19 /G

If you wear dentures, make sure they fit properly. Avoiding sources of oral irritation (such as dentures that don't fit properly) may help lower your risk for oral cancer.

Hyperlinks

- 1. www.cancer.org/cancer/risk-prevention/tobacco.html
- 2. www.cancer.org/cancer/types/cervical-cancer.html
- 3. www.cancer.org/cancer/risk-prevention/hpv.html
- 4. www.cancer.org/cancer/risk-prevention/sun-and-uv/uv-radiation.html
- 5. www.cancer.org/cancer/types/skin-cancer.html
- 6. <u>www.cancer.org/cancer/risk-prevention/diet-physical-activity/acs-guidelines-nutrition-physical-activity-cancer-prevention.html</u>
- 7. <u>www.cancer.org/cancer/types/oral-cavity-and-oropharyngeal-cancer/about/what-is-oral-cavity-cancer.html</u>

References

Chainani-Wu N. Diet and oral, pharyngeal, and esophageal cancer. *Nutr Cancer*. 2002;44(2):104-126. doi:10.1207/S15327914NC4402_01.

Chaturvedi AK, D'Souza G, Gillison ML, Katki HA. Burden of HPV-positive oropharynx cancers among ever and never smokers in the U.S. population. *Oral Oncol.* 2016;60:61-67. doi:10.1016/j.oraloncology.2016.06.006.

Leeman JE, Katabi N, Wong RJ, Lee NY, Romesser PB. Ch. 65 - Cancer of the Head and Neck. In: Niederhuber JE, Armitage JO, Doroshow JH, Kastan MB, Tepper JE, eds. *Abeloff's Clinical Oncology*. 6th ed. Philadelphia, Pa. Elsevier; 2020.

Lucenteforte E, Garavello W, Bosetti C, La Vecchia C. Dietary factors and oral and pharyngeal cancer risk. *Oral Oncol.* 2009;45(6):461-467. doi:10.1016/j.oraloncology.2008.09.002.

National Cancer Institute. Physician Data Query (PDQ). Lip and Oral Cavity Cancer Treatment (Adult) (PDQ®)—Health Professional Version. September 05, 2019. Accessed at https://www.cancer.gov/types/head-and-neck/hp/adult/lip-mouth-treatment-pdq on September 29, 2020.

Rock CL, Thomson C, Gansler T, et al. American Cancer Society guideline for diet and physical activity for cancer prevention. *CA: A Cancer Journal for Clinicians*. 2020;70(4). doi:10.3322/caac.21591. Accessed at https://onlinelibrary.wiley.com/doi/full/10.3322/caac.21591 on June 9, 2020.

Saslow D, Andrews KS, Manassaram-Baptiste D, et al. Human papillomavirus vaccination 2020 guideline update: American Cancer Society guideline adaptation. *CA Cancer J Clin.* 2020; DOI: 10.3322/caac.21616.

Tumban E. A Current Update on Human Papillomavirus-Associated Head and Neck Cancers. *Viruses*. 2019;11(10):922. Published 2019 Oct 9. doi:10.3390/v11100922.

Last Revised: March 23, 2021

Written by

The American Cancer Society medical and editorial content team (https://www.cancer.org/cancer/acs-medical-content-and-news-staff.html)

Our team is made up of doctors and oncology certified nurses with deep knowledge of cancer care as well as journalists, editors, and translators with extensive experience in medical writing.

American Cancer Society medical information is copyrighted material. For reprint requests, please see our Content Usage Policy (www.cancer.org/about-us/policies/content-usage.html).

cancer.org | 1.800.227.2345